PCT/EP2005/001887

21

CLAIMS

- 1. An apparatus for fetal monitoring comprising:
- 5 a) means for determining a fetal heart rate,
 - b) means for identifying a primary fetal heart rate component which is required to shift a volume of blood from the heart to the cardiovascular system,
 - c) means for subtracting the primary component from the
- 10 determined fetal heart rate to determine a residual component; and
 - d) means for using said residual component for analysis of the fetal heart rate beat-to-beat variation.
- 2. An apparatus as claimed in claim 1, wherein the primary fetal heart rate component is identified through a polynomial curve fit approximation of the fetal heart rate data.
- 3. An apparatus as claimed in claim 1, wherein said means for identifying the primary fetal heart rate component is adapted to perform the following steps:
 - i) linear interpolation of recorded fetal heart rate data;
- 25 ii) resampling at a resampling frequency, thereby forming a resampled series of fetal heart rate data, and; iii) polynomial curve fit approximation of said resampled series.
- 30 4. An apparatus as claimed in claim 2 or 3, wherein the polynomial curve fit approximation utilises polynomials of at least the 5^{th} order.

WO 2005/079666 PCT/EP2005/001887

22

5. An apparatus as claimed in claim 4, wherein said polynomials are of the $5^{\rm th}$ order.

- 6. An apparatus as claimed in claim 4, wherein said polynomials are of the 12th order.
 - 7. An apparatus as claimed in any one of claim 2 to 6, wherein the polynomial approximation is obtained through a least squares approximation.

10

- 8. An apparatus as claimed in any preceding claim, wherein the primary fetal heart rate component is determined by:
- i) dividing the fetal heart rate data into regionsof a predetermined size; and
 - ii) performing individual polynomial approximations in each region.
- 9. An apparatus as claimed in claim 8, wherein each 20 polynomial approximation is constrained such that neighbouring polynomial functions align and have equal first derivatives at the region border where they join.
- 10. An apparatus as claimed in claim 8 or 9, wherein the predetermined size is greater than or equal to 20 consecutive heart rate samples.
 - 11. An apparatus as claimed in claim 10, wherein the predetermined size is 20 consecutive heart rate samples.

30

12. An apparatus as claimed in any preceding claim, wherein the means for using said residual component for analysis of the fetal heart rate beat-to-beat variation

PCT/EP2005/001887

WO 2005/079666

25

is adapted to apply statistical tests for analysing the residual component in order to determine the response of the fetus.

- 13. An apparatus as claimed in claim 12, wherein the statistical test comprises monitoring of a 95th percentile of the fetal heart rate residual component.
- 14. An apparatus as claimed in claim 13, wherein the statistical test further comprises calculating a median and a variance of said 95th percentile over a predetermined period of time.
- 15. An apparatus as claimed in claim 14, wherein saidpredetermined period of time is longer than 10 minutes.
- 16. An apparatus as claimed in any one of claims 13 15, wherein if the median of the 95th percentile is consistently below 3ms the fetal heart rate is classed as abnormal and non-reactive.
 - 17. An apparatus as claimed in any one of claims 12 15, wherein said means for using said residual component for analysis of the fetal heart rate beat-to-beat variation is adapted to indicate a significant reduction of fetal reactivity given a recording of the median of the 95th percentile below 2.3 ms and the variance of the 95th percentile below 0.1 over an extended period of time.
- 30 18. An apparatus as claimed in any one of claims 12 15, wherein said means for using said residual component for analysis of the fetal heart rate beat-to-beat variation is adapted to indicate a significant reduction of fetal

WO 2005/079666 PCT/EP2005/001887

24

reactivity given a recording of a decreasing trend of the median of the $95^{\rm th}$ percentile over an extended period of time.

- 5 19. An apparatus as claimed in any one of claims 12 15, wherein said means for using said residual component for analysis of the fetal heart rate beat-to-beat variation is adapted to exclude an abnormally low fetal heart rate variation if the median of the 95th percentile is consistently above 3ms.
 - 20. An apparatus as claimed in claim 12, wherein the statistical test comprises monitoring of a short term, e.g. 3-4ms, frequency distribution of the fetal heart rate residual component.
 - 21. An apparatus as claimed in claim 20, wherein if a 3-4ms frequency distribution is less than 7% the fetal heart rate is classed as non-reactive.

20

15

- 22. A method for fetal monitoring comprising the steps of:
- a) determining a fetal heart rate,
- b) identifying a primary fetal heart rate component which
- 25 is required to shift a volume of blood from the heart to the cardiovascular system,
 - c) subtracting the primary component from the determined fetal heart rate to determine a residual component; and
 - d) using said residual component for analysis of the
- 30 fetal heart rate beat-to-beat variation.
 - 23. A computer program for executing the steps of:
 - a) determining a fetal heart rate,

WO 2005/079666 PCT/EP2005/001887

25

- b) identifying a primary fetal heart rate component which is required to shift a volume of blood from the heart to the cardiovascular system,
- c) subtracting the primary component from the determined fetal heart rate to determine a residual component; and d) using said residual component for analysis of the fetal heart rate beat-to-beat variation.